

	<p style="text-align: center;">IPU</p> <p style="text-align: center;">Hospital</p> <p style="text-align: center;">Community</p> <p style="text-align: center;">Other (please state):</p> <p style="text-align: center;">.....</p>
Age	
Sex	
Diagnosis	
Reason for SC PPI	
Switch from oral PPI? (please circle)	<p>Yes</p> <p>No</p>
SC PPI chosen (please circle)	<p>Esomeprazole</p> <p>Omeprazole</p> <p>Pantoprazole</p>
Dose of SC PPI (mg)	
Reason for choice of SC PPI (e.g. availability, familiarity, ease of administration)	
SC PPI administration method chosen (please circle)	<p>CSCI</p> <p>SC infusion</p> <p>SC bolus</p>

Diluent used (please circle)	<p style="text-align: center;">0.9% sodium chloride</p> <p style="text-align: center;">Water for injection</p>
Duration of administration (minutes/hours)	
Length of time on SC PPI (days)	
Reason for stopping SC PPI (e.g. no symptomatic benefit, clinical deterioration, death etc.)	
Renal function if known (numerical/NA)	
Evidence of symptomatic benefit (e.g. reduced pain, reduced nausea etc.)	
Adverse events (please circle)	<p style="text-align: center;">None</p> <p style="text-align: center;">Site irritation – mild</p> <p style="text-align: center;">Site irritation – moderate</p> <p style="text-align: center;">Site irritation – severe</p> <p style="text-align: center;">Clinically significant renal impairment</p> <p style="text-align: center;">Clostridium difficile infection</p> <p style="text-align: center;">Hypomagnesaemia</p> <p style="text-align: center;">Hyponatraemia</p> <p style="text-align: center;">Other (please state):</p> <p style="text-align: center;">.....</p>
Comments (please add any additional comments below)	